

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
5 July 2007 (05.07.2007)

PCT

(10) International Publication Number
WO 2007/075621 A1

(51) International Patent Classification:

A61K 31/05 (2006.01) A61K 31/375 (2006.01)

A61K 31/355 (2006.01) A61K 31/436 (2006.01)

(21) International Application Number:

PCT/US2006/048329

(22) International Filing Date:

19 December 2006 (19.12.2006)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

60/752,189 20 December 2005 (20.12.2005) US

(71) Applicant (for all designated States except US): **WYETH**
[US/US]; Five Giralda Farms, Madison, NJ 07940 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **RUBINO, Joseph, Thomas** [US/US]; 4 Sunrise Way, Towaco, NJ 07082 (US). **POOJA, Gandhi** [US/US]; 1 Ash Court, Highland Mills, NY 10930 (US). **PHELAN, Lynn** [US/US]; 15 Sheffield Street, Lake Hiawatha, NJ 07034 (US).

(74) Agents: **KODROFF, Cathy, A.** et al.; Howson And Howson, 501 Office Center Drive, Suite 210, Fort Washington, PA 19034 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: CONTROL OF CCI-779 DOSAGE FORM STABILITY THROUGH CONTROL OF DRUG SUBSTANCE IMPURITIES

(57) Abstract: A method of preparing a rapamycin composition having increased potency is provided. The method involves selecting a rapamycin compound having less than 1.5% oxidative and hydrolytic rapamycin impurities and formulating the selected rapamycin with an antioxidant and optional excipients.

WO 2007/075621 A1

CONTROL OF CCI-779 DOSAGE FORM STABILITY THROUGH CONTROL OF DRUG SUBSTANCE IMPURITIES

5 BACKGROUND OF THE INVENTION

Rapamycin 42-ester with 3-hydroxy-2-(hydroxymethyl)-2-methylpropionic acid (CCI-779) has potential as an antitumor agent. This compound is now known generically under the name temsirolimus (Wyeth). The preparation and use of hydroxyesters of rapamycin, including temsirolimus, are described in US Patent Nos.
10 5,362,718 and 6,277,983.

The intravenous dosage form includes a solution of CCI-779, α -tocopherol, citric acid, and ethanol in propylene glycol. Rapamycin and related compounds may be susceptible to oxidative and hydrolytic degradation during synthesis and purification of the drug substance or when formulated as a dosage form. Oxidation
15 generally begins via peroxidation of unsaturated carbons in the 1-7 carbon polyene region of rapamycin and its derivatives, such as CCI-779. The initial peroxidation generally proceeds to form a number of oxide, hydroxide and aldehyde degradation products.

Collectively, these degradation products or impurities are referred to as "group
20 II" or "oxidative and hydrolytic" degradation products or impurities. The presence of these impurities/degradation products can catalyze degradation of the drug and thereby destabilize the drug when present in sufficiently high amounts.

Addition of antioxidants to both the drug substance, during processing, and in the final drug product may inhibit degradation caused by the degradation products or
25 impurities. However, when these degradation/impurity levels reach a critical value, further degradation of drug product is difficult to inhibit by practical means. This is especially a limitation for parenteral products because the levels of antioxidants and other stabilizers used in a formulation is often limited by safety concerns and their levels in new products may be limited by previous human safety experience.

30 Because of the negative influence of oxidative/hydrolytic degradation products on the potency and purity of drug product, it is advantageous to limit their amount in the composition of the final drug product.

US Patent No. 6,605,613 B2 discusses stabilization of macrolides using various antioxidants. The primary focus of that patent is to stabilize drug during its preparation and final isolation.

Because of the variety of degradation pathways that occur during oxidation
5 and hydrolysis of rapamycin derivatives as well as the ability of rapamycin and related compounds to form various isomers, the isolation and quantitation of oxidative/hydrolytic degradation products/impurities as individual substances is difficult to achieve. For this reason, it is often necessary to quantitate
oxidative/hydrolytic degradation products as a group rather than as single, individual
10 compounds.

What is needed in the art are alternate methods for preparing rapamycin compositions that have less degradation impurities.

SUMMARY OF THE INVENTION

15 In one aspect, the present invention provides a method of preparing a rapamycin composition having increased potency.

In another aspect, the present invention provides a method for preparing a rapamycin composition having increased potency by formulating a rapamycin compound having not more than 1.5% oxidative and hydrolytic rapamycin impurities
20 with an antioxidant and optional excipients.

Other aspects and advantages of the invention will be readily apparent from the following detailed description of the invention.

BRIEF DESCRIPTION OF THE DRAWING

25 Fig. 1 is an LC/UV chromatogram of an exemplary temsirolimus (CCI-779) sample preparation which is corrected for solvent.

Fig. 2 is an LC/MS chromatogram of an exemplary temsirolimus (CCI-779) sample preparation. More particularly, this HPLC/MS Chromatogram for oxidative / hydrolysis degradants has a time in column (TIC) Range: m/z 1044.7 to 1076.7.


30 Fig. 3 is a plot of total nonoxidative degradants and oxidative/hydrolysis degradants vs. time for a parenteral drug product of temsirolimus (CCI-779) prepared with drug substance that contained 0.5, 1, or 2% initial oxidative/hydrolysis

degradants. In this figure, "OD" refers to the percent (%) oxidative/hydrolysis degradants initially in the drug substance.

DETAILED DESCRIPTION OF THE INVENTION

5 The present invention provides methods of preparing rapamycin compositions having increased stability. The term "increased stability" as used herein refers to a rapamycin composition whereby the concentration of the rapamycin compound contained therein decreases to a lesser extent over time and has fewer or lower levels of degradation products as compared to rapamycin compositions in the art. Desirably,
10 the rapamycin compositions of the present invention show minimal degradation after storage at 25 °C or 40 °C compared to compositions where the oxidative/hydrolytic impurities in the starting material are not controlled.

 As used herein, the term rapamycin compound defines a class of immunosuppressive compounds that contain the basic rapamycin nucleus as shown
15 below.



 The rapamycin compounds of this invention include compounds that are
20 chemically or biologically modified as derivatives of the rapamycin nucleus, while still retaining immunosuppressive properties. Accordingly, the term rapamycin compound includes rapamycin, and esters, ethers, carbamates, oximes, hydrazones, and hydroxylamines of rapamycin, as well as rapamycins in which functional groups on the rapamycin nucleus have been modified, for example through reduction or
25 oxidation.

 The term rapamycin compound also includes 42- and/or 31-esters and ethers of rapamycin as described in the following patents, which are all hereby incorporated

by reference: alkyl esters (US Patent No. 4,316,885); aminoalkyl esters (US Patent No. 4,650,803); fluorinated esters (US Patent No. 5,100,883); amide esters (US Patent No. 5,118,677); carbamate esters (US Patent No. 5,118, 678); silyl esters (US Patent No. 5,120,842); aminodiester (US Patent No. 5,162,333); sulfonate and sulfate esters (US Patent No. 5,177,203); esters (US Patent No. 5,221,670); alkoxyesters (US Patent No. 5,233,036); O-aryl, -alkyl, -alkenyl, and -alkynyl ethers (US Patent No. 5,258,389); carbonate esters (US Patent No. 5,260,300); arylcarbonyl and alkoxy carbonyl carbamates (US Patent No. 5,262,423); carbamates (US Patent No. 5,302,584); hydroxyesters (US Patent No. 5,362,718); hindered esters (US Patent No. 5,385,908); heterocyclic esters (US Patent No. 5,385,909); gem-disubstituted esters (US Patent No. 5,385,910); amino alkanolic esters (US Patent No. 5,389,639); phosphorylcarbamate esters (US Patent No. 5,391,730); carbamate esters (US Patent No. 5,411,967); carbamate esters (US Patent No. 5,434,260); amidino carbamate esters (US Patent No. 5,463,048); carbamate esters (US Patent No. 5,480,988); carbamate esters (US Patent No. 5,480,989); carbamate esters (US Patent No. 5,489,680); hindered N-oxide esters (US Patent No. 5,491,231); biotin esters (US Patent No. 5,504,091); O-alkyl ethers (US Patent No. 5,665,772); and PEG esters of rapamycin (US Patent No. 5,780,462). The preparation of these esters and ethers is disclosed in the patents listed above.

Further included within the definition of the term rapamycin compound are 27-esters and ethers of rapamycin, which are discussed in US Patent No. 5,256,790. Also described are C-27 ketone rapamycins which are reduced to the corresponding alcohol, which is in turn converted to the corresponding ester or ether. The preparation of these esters and ethers is discussed in the patents provided above. Also included are oximes, hydrazones, and hydroxylamines of rapamycin as discussed in US Patent Nos. 5,373,014; 5,378,836; 5,023,264; and 5,563,145. The preparation of these oximes, hydrazones, and hydroxylamines is discussed in the above-listed patents. The preparation of 42-oxorapamycin is discussed in US Patent No. 5,023,263.

The term rapamycin compound also refers to any combination of different rapamycins or chemical compounds that contains rapamycin or any derivative thereof.

Specific examples of rapamycin compounds that can be used in the invention include, without limitation, rapamycin, CCI-779, norrapamycin, deoxorapamycin, desmethylrapamycin, desmethoxyrapamycin, or the rapamycins described in US Patent Publication No. 2006-0135549 (claiming priority from US Provisional Application No. 60/637,666) and US Patent Publication No. 2006-013550 A1, (claiming priority from US Provisional Application No. 60/638,004), which are hereby incorporated by reference, or pharmaceutically acceptable salts, prodrugs, or metabolites thereof, among others.

The term “desmethylrapamycin” refers to the class of rapamycin compounds which lack one or more methyl groups. Examples of desmethylrapamycins that can be used according to the present invention include 3-desmethylrapamycin (US Patent No. 6,358,969), 7-O-desmethyl-rapamycin (US Patent No. 6,399,626), 17-desmethylrapamycin (US Patent No. 6,670,168), and 32-O-desmethylrapamycin, among others.

The term “desmethoxyrapamycin” refers to the class of rapamycin compounds which lack one or more methoxy groups and includes, without limitation, 32-desmethoxyrapamycin.

The rapamycin compositions of the invention include the rapamycin compound at an amount sufficient to treat the conditions and diseases identified below. Specifically, the rapamycin compound is present in the rapamycin compositions at about 0.1 to 30 wt%, 0.5 to 25 wt%, 1 to 20 wt%, 5 to 15 wt%, or 7 to 12 wt% (wt/wt). Desirably, the rapamycin compound is present at an amount of 2 to about 500 mg, 5 mg to 250 mg, 10 mg to 100 mg, 15 mg to 50 mg, or about 20 mg to 25 mg.


The rapamycin compound of the invention can be in a micronized or nonmicronized form and can also include tautomeric forms of the rapamycin compound. The present invention also includes derivatives of rapamycin, including, but not limited to, esters, carbamates, sulfates, ethers, oximes, carbonates, and the like.

The rapamycin compound can also encompass “metabolites” which are unique products formed by processing rapamycin by the cell or patient. Desirably, metabolites are formed *in vivo*.

It is also desirable that the rapamycin compound in the compositions of the invention degrades less than the rapamycin compound in the compositions in the art. Of course, it is most desirable that the concentration of the rapamycin compound in the compositions of the present invention be maintained. However, it is desirable that the concentration of the rapamycin compound in the compositions of the invention degrades less than about 2 % after storage for 3-5 months at 25 °C or 1 month at 40 °C, more desirably less than about 1 %.

The inventors found that more potent rapamycin compositions are obtained when the rapamycin compound utilized therein contained less than 1.5% (*i.e.*, 0 or 0.01, 0.01 to 1.5%) oxidative and hydrolytic impurities in the starting material. In fact, by utilizing rapamycin compounds containing 0.5 to 1% or less oxidative and hydrolytic impurities, degradation of the rapamycin was significantly reduced or eliminated. More desirably, the rapamycin compound contains less than about 1%, less than about 0.5%, less than about 0.4%, less than about 0.3%, less than about 0.2%, or about 0.1% oxidative impurities. Most desirably, the rapamycin contains less than about 0.5% oxidative impurities.

The term "oxidative and hydrolytic impurities of rapamycin", or variations thereof as used herein, refers to chemical compounds that form in rapamycin compositions. These impurities can include a group of oxygen addition compounds involving the C₁₋₆ region of rapamycin or its analogs identified below. These impurities can therefore include aldehydes, epoxides, hydroxides, and combinations thereof of rapamycin or rapamycin derivatives. These impurities can also include ring-opened forms of rapamycin or rapamycin analogs that contain the oxygen addition modifications described above for the C₁₋₆ region.



The presence of these oxidative and hydrolytic impurities is typically measured using high performance liquid chromatography (HPLC) with ultraviolet

(UV) or mass spectrometric (MS) detection. Specifically, the oxidative and degradation impurities can be quantitated using either HPLC/UV or HPLC/MS. More specifically, the oxidative and hydrolytic impurities may be quantitated as a mixture of co-eluting materials over a specified range of retention times (HPLC/UV). For example, in one embodiment, the retention time of the CCI-779 Isomer B peak should be between 18 and 24 minutes using a suitable chromatograph column (*e.g.*, a reverse phase column). Alternatively quantitation by analyzing the extent of one oxygen, two oxygen, 3 oxygen, one oxygen plus water, and water incorporation, based on the m/z of the addition product, is employed.

10 The method of the invention thereby includes preparing a rapamycin composition by selecting a rapamycin compound as noted above for use therein. Desirably, the rapamycin compound has less than 1.5% oxidative and hydrolytic rapamycin impurities. After selection of the desired rapamycin compound, it is formulated with one or more of an antioxidant.

15 Antioxidants that can be used in the rapamycin compositions of the present invention include, but are not limited to, citric acid, alpha tocopherol, BHA, BHT (2,6-di-*tert*-butyl-4-methylphenol), monothioglycerol, Vitamin C, and propyl gallate. In one embodiment, Vitamin C is ascorbic acid. However, one of skill in the art may substitute a pharmaceutically acceptable salt thereof for the ascorbic acid. Desirably, the antioxidant is d,l- α -tocopherol. In one embodiment, the antioxidant may be used in concentrations ranging from 0.0005 wt% to 3 wt%, and desirably from 0.001 wt% to 3 wt%.

25 The rapamycin compositions of the invention may also contain suitable excipients including, without limitation, water soluble polymers, pH modifying agents, chelating agents, surfactants, fillers, binders, disintegrants, and the like. Any given rapamycin composition useful in the invention may contain multiple ingredients of each class of component. For example, some compositions may contain one or more antioxidant.

30 pH modifying agents include, but are not limited to, citric acid, sodium citrate, acetic acid, lactic acid, dilute HCl, and other mild acids or bases capable of buffering a solution containing the rapamycin compound to a pH in the range of about 4 to about 6.

Chelating agents, and other materials capable of binding metal ions, can be included in the rapamycin compositions of the invention. Desirably, the chelating agent enhances the stability of the rapamycin compound. In certain embodiments, the antioxidant component of the formulation of the invention can exhibit chelating activity. Examples of chelating agents include, without limitation, citric acid and ascorbic acid (which may function as both a classic antioxidant and a chelating agent in the present formulations). Other chelating agents include such materials as are capable of binding metal ions in solution, such as ethylene diamine tetra acetic acid (EDTA), its salts, or amino acids such as glycine, which are capable of enhancing the stability of the rapamycin compound. Typically, chelating agents are used in the lower end of the range of concentrations for the antioxidant component provided herein. In one example, citric acid is utilized at a concentration of less than 0.01% w/v. Additionally, such chelating agents may be used in combination with other antioxidants as part of the antioxidant component of the invention. For example, an acceptable formulation may contain both citric acid and d,l- α -tocopherol. Optimal concentrations for the selected antioxidant(s) can be readily determined by one of skill in the art, based upon the information provided herein.

Surfactants may include polysorbate 80, polyoxyethylene fatty acid esters, sodium lauryl sulfate, sodium dodecyl sulfate, salts of bile acids (taurocholate, glycocholate, cholate, deoxycholate, etc.) that may be combined with lecithin, Vitamin E TPGS, and/or poloxamers. The surfactant can be present in the rapamycin compositions at 0.5 to 10 wt%, 1 to 8 wt%, or 3 to 5 wt% (wt/wt), or can be present in amounts from the lower or higher end of these ranges up to about 50 wt%.

Binders, fillers, and disintegrants can include sucrose, lactose, microcrystalline cellulose, croscarmellose sodium, magnesium stearate, gum acacia, cholesterol, tragacanth, stearic acid, gelatin, casein, lecithin (phosphatides), carboxymethylcellulose calcium, carboxymethylcellulose sodium, methylcellulose, hydroxyethylcellulose, hydroxypropylcellulose, hydroxypropylmethylcellulose phthalate, noncrystalline cellulose, cetostearyl alcohol, cetyl alcohol, cetyl esters wax, dextrates, dextrin, lactose, dextrose, glyceryl monooleate, glyceryl monostearate, glyceryl palmitostearate, polyoxyethylene alkyl ethers, polyethylene glycols,

polyoxyethylene castor oil derivatives, polyoxyethylene stearates, and polyvinyl alcohol.

Typical water soluble polymers include, but are not limited to, polyvinylpyrrolidone (PVP), hydroxypropylmethylcellulose (HPMC), polyethylene glycol (PEG), and cyclodextrin or mixtures thereof. Desirably, the water-soluble polymer is PVP and has a molecular weight of between 2.5 and 60 kilodaltons.

The rapamycin compositions described herein can be formulated in any form suitable for the desired route of delivery using a pharmaceutically effective amount of the rapamycin compound. For example, the compositions of the invention can be delivered by a route such as oral, dermal, transdermal, intrabronchial, intranasal, intravenous, intramuscular, subcutaneous, parenteral, intraperitoneal, intranasal, vaginal, rectal, sublingual, intracranial, epidural, intratracheal, or by sustained release.

Suitable oral formulations for the rapamycin compositions can be prepared as described for CCI-779, as described in International Patent Publication No. WO 2004/026280 and US Patent Application Publication No. US 2004-0077677 A1, which are hereby incorporated by reference. In one embodiment, the composition contains 0.1 to 30 wt%, 0.5 to 25 wt%, 1 to 20 wt%, 5 to 15 wt%, or 7 to 12 wt% (wt/wt) of a rapamycin compound and 0.001 wt% to 1 wt%, 0.01 wt% to 1 wt%, or 0.1 wt% to 0.5 wt% (wt/wt) of an antioxidant. The compositions can optionally contain 0.5 to 50 wt%, 1 to 40 wt%, 5 to 35 wt%, 10 to 25 wt%, or 15 to 20 wt% (wt/wt) of a water soluble polymer and 0.5 to 10 wt%, 1 to 8 wt%, or 3 to 5 wt% (wt/wt) of a surfactant. However, other embodiments may contain more, or less, of these components.

Oral formulations may include any conventionally used oral forms, including tablets, capsules, buccal forms, troches, lozenges and oral liquids, suspensions or solutions. Capsules may contain mixtures of the rapamycin compound with inert fillers and/or diluents such as the pharmaceutically acceptable starches (e.g. corn, potato or tapioca starch), sugars, artificial sweetening agents, powdered celluloses, such as crystalline and microcrystalline celluloses, flours, gelatins, gums, etc. Useful tablet formulations may be made by conventional compression, wet granulation or dry granulation methods and utilize pharmaceutically acceptable diluents, binding agents, lubricants, disintegrants, surface modifying agents (including

surfactants), suspending or stabilizing agents, including, but not limited to, magnesium stearate, stearic acid, talc, sodium lauryl sulfate, microcrystalline cellulose, carboxymethylcellulose calcium, polyvinylpyrrolidone, gelatin, alginic acid, acacia gum, xanthan gum, sodium citrate, complex silicates, calcium carbonate, glycine, dextrin, sucrose, sorbitol, dicalcium phosphate, calcium sulfate, lactose, kaolin, mannitol, sodium chloride, talc, dry starches and powdered sugar. Surface modifying agents can include nonionic and anionic surface modifying agents. Representative examples of surface modifying agents include, but are not limited to, poloxamer 188, benzalkonium chloride, calcium stearate, cetostearyl alcohol, cetomacrogol emulsifying wax, sorbitan esters, colloidal silicon dioxide, phosphates, sodium dodecylsulfate, magnesium aluminum silicate, and triethanolamine. Oral formulations herein may utilize standard delay or time release formulations to alter the absorption of the rapamycin. The oral formulation may also include water or a fruit juice, containing appropriate solubilizers or emulsifiers as needed.

In some cases it may be desirable to administer the rapamycin composition directly to the airways in the form of an aerosol.

The rapamycin compositions may also be administered parenterally or intraperitoneally. Solutions or suspensions of the rapamycin compound as a free base or pharmacologically acceptable salt can be prepared in water suitably mixed with a surfactant such as hydroxy-propylcellulose. Dispersions can also be prepared in glycerol, liquid polyethylene glycols and mixtures thereof in oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms.

The pharmaceutical forms suitable for injectable use include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. In all cases, the form must be sterile and must be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (e.g., glycerol, propylene glycol and liquid polyethylene glycol), suitable mixtures thereof, and vegetable oils.

Particularly suitable injectable formulations containing the rapamycin compound can be prepared in a manner similar to those described in International Patent Publication No. WO 2004/011000 and US Patent Application Publication No. US 2004-0167152 A1, which are hereby incorporated by reference. In this

5 embodiment, the injectable formulation useful in the invention provides a rapamycin composition cosolvent concentrate containing a parenterally acceptable solvent and an antioxidant as described above and a parenteral formulation containing a rapamycin compound, a parenterally acceptable cosolvent, an antioxidant, a diluent solvent, and a surfactant. For example, a parenterally acceptable solvent can include a non-

10 alcoholic solvent, an alcoholic solvent, or mixtures thereof. Examples of suitable non-alcoholic solvents include, without limitation, dimethylacetamide, dimethylsulfoxide, or mixtures thereof. Examples of alcoholic solvent include, without limitation, one or more alcohols as the alcoholic solvent component of the formulation. Examples of solvents useful in the formulations invention include,

15 without limitation, ethanol, propylene glycol, polyethylene glycol 300, polyethylene glycol 400, polyethylene glycol 600, polyethylene glycol 1000, or mixtures thereof. Further, ethanol and propylene glycol can be combined to produce a less flammable product. These latter two cosolvents are particularly desirable because degradation via oxidation and lactone cleavage occurs to a lower extent for these cosolvents. Larger

20 amounts of ethanol in the mixture generally result in better chemical stability. A concentration of 30 to 100% v/v of ethanol in the mixture is desirable.

The stability of the rapamycin compound in the parenterally acceptable alcoholic cosolvent can be enhanced by addition of an antioxidant to the formulation. Generally, the parenteral formulation useful in the invention will contain an

25 antioxidant component(s) in a concentration of 0.001% to 3% w/v or 0.01% to 0.1% w/v, of the cosolvent concentrate, although lower or higher concentrations may be desired. Of the antioxidants, d,l- α -tocopherol is particularly desirable and is used at a concentration of 0.01 to 0.1% w/v with a concentration of 0.075% w/v of the cosolvent concentrate being most desirable.

30 Dosage regimens are expected to vary according to the route of administration. For example, dosages for oral administration are often up to five to tenfold greater than for intravenous (i.v.) administration. In one embodiment, a dosage of the

rapamycin compound may be about 2 to about 500 mg/day, 5 mg/day to 75 mg/day, 10 mg/day to 50 mg/day, 15 mg/day to 35 mg/day, or about 20 mg/day to 25 mg/day for an adult. However, this dosage can be adjusted upwardly or downwardly by one of skill in the art, depending upon the indication being treated, the size of the patient, and other factors which are known those of skill in the art.

In certain embodiments of the parenteral formulations useful in the invention, precipitation of the rapamycin compound upon dilution with aqueous infusion solutions or blood is prevented through the use of a surfactant contained in the diluent solution. The most important component of the diluent is a parenterally acceptable surfactant. One particularly desirable surfactant is polysorbate 20 or polysorbate 80. However, one of skill in the art may readily select other suitable surfactants from among salts of bile acids (taurocholate, glycocholate, cholate, deoxycholate, etc.) which are optionally combined with lecithin. Alternatively, ethoxylated vegetable oils, such as a pegylated castor oil (e.g., such as PEG-35 castor oil which is sold, e.g., under the name Cremophor EL, BASF), vitamin E tocopherol propylene glycol succinate (Vitamin E TGPS), and polyoxyethylene-polyoxypropylene block copolymers can be used in the diluent as a surfactant, as well as other members of the polysorbate family such as polysorbate 20 or 60. Other components of the diluent may include water, ethanol, polyethylene glycol 300, polyethylene 400, polyethylene 600, polyethylene 1000, or blends containing one or more of these polyethylene glycols, propylene glycol and other parenterally acceptable cosolvents or agents to adjust solution osmolarity such as sodium chloride, lactose, mannitol or other parenterally acceptable sugars, polyols and electrolytes. It is expected that the surfactant will include at least 5% w/v, at least 10% w/v, or at least 5% w/v of the diluent solution. Desirably, the surfactant will include 2 to 100% w/v, 5 to 80% w/v, 10 to 75% w/v, or 15 to 60 % w/v of the diluent solution.

A parenteral formulation useful in the invention can be prepared as a single solution, or desirably can be prepared as a cosolvent concentrate containing the rapamycin compound, an alcoholic solvent, and an antioxidant, which is subsequently combined with a diluent that contains a diluent solvent and suitable surfactant. Prior to use, the cosolvent concentrate is mixed with a diluent comprising a diluent solvent, and a surfactant. When rapamycin compound is prepared as a cosolvent concentrate

according to this invention, the concentrate can contain concentrations of the rapamycin compound from 0.05 mg/mL, from 2.5 mg/mL, from 5 mg/mL, from 10 mg/mL or from 25 mg/mL up to approximately 50 mg/mL. The concentrate can be mixed with the diluent up to approximately 1 part concentrate to 1 part diluent, to give
5 parenteral formulations having concentrations of the rapamycin compound from 1 mg/mL, from 5 mg/mL, from 10 mg/mL, from 20 mg/mL, up to approximately 25 mg/mL. For example the concentration of the rapamycin compound in the parenteral formulation may be from about 2.5 to 10 mg/mL. This invention also covers the use of formulations having lesser concentrations of the rapamycin compound in the
10 cosolvent concentrate, and formulations in which one part of the concentrate is mixed with greater than 1 part of the diluent, *e.g.*, concentrate: diluent in a ratio of about 1:1.5, 1:2, 1:3, 1:4, 1:5, or 1:9 v/v and so on, to rapamycin compound parenteral formulations having a rapamycin compound concentration down to the lowest levels of detection. Typically the antioxidant may include from about 0.0005 to 0.5% w/v of
15 the formulation. The surfactant may for example include from about 0.5 % to about 10% w/v of the formulation. The alcoholic solvent may for example include from about 10% to about 90% w/v of the formulation.

The parenteral formulations useful in this invention can be used to produce a dosage form that is suitable for administration by either direct injection or by addition
20 to sterile infusion fluids for intravenous infusion.

Transdermal administrations include all administrations across the surface of the body and the inner linings of bodily passages including epithelial and mucosal tissues. Such administrations may be carried out using the present compounds, or pharmaceutically acceptable salts thereof, in lotions, creams, foams, patches,
25 suspensions, solutions, and suppositories (rectal and vaginal). Transdermal administration may be accomplished through the use of a transdermal patch containing the active compound and a carrier that is inert to the active compound, is non toxic to the skin, and allows delivery of the agent for systemic absorption into the blood stream via the skin. The carrier may take any number of forms such as creams
30 and ointments, pastes, gels, and occlusive devices. The creams and ointments may be viscous liquid or semisolid emulsions of either the oil-in-water or water-in-oil type. Pastes comprised of absorptive powders dispersed in petroleum or hydrophilic

petroleum containing the active ingredient may also be suitable. A variety of occlusive devices may be used to release the active ingredient into the blood stream such as a semi-permeable membrane covering a reservoir containing the active ingredient with or without a carrier, or a matrix containing the active ingredient.

5 Other occlusive devices are known in the literature.

Suppository formulations may be made from traditional materials, including cocoa butter, with or without the addition of waxes to alter the suppository's melting point, and glycerin. Water soluble suppository bases, such as polyethylene glycols of various molecular weights, may also be used.

10 The rapamycin compound of the invention may be formulated for any suitable delivery route and vehicle and assembled in the form of a kit of parts.

Thus, the rapamycin compositions of the invention can be useful as an antineoplastic agent, and therefore in the treatment of solid tumors, including sarcomas and carcinomas; and more particularly against astrocytomas, prostate cancer, breast cancer, colon cancer, small cell lung cancer, and ovarian cancer; and adult T-cell leukemia/lymphoma. The rapamycin compound-containing compositions are also useful in the treatment or inhibition of transplantation rejection such as kidney, heart, liver, lung, bone marrow, pancreas (islet cells), cornea, small bowel, and skin allografts, and heart valve xenografts; in the treatment or inhibition of graft vs. host disease; in the treatment or inhibition of autoimmune diseases such as lupus, rheumatoid arthritis, diabetes mellitus, myasthenia gravis, and multiple sclerosis; and diseases of inflammation such as psoriasis, dermatitis, eczema, seborrhea, inflammatory bowel disease, pulmonary inflammation (including asthma, chronic obstructive pulmonary disease, emphysema, acute respiratory distress syndrome, bronchitis, and the like) and ocular uveitis; adult T-cell leukemia/lymphoma; fungal infections; hyperproliferative vascular diseases such as restenosis; graft vascular atherosclerosis; and cardiovascular disease, cerebral vascular disease, and peripheral vascular disease, such as coronary artery disease, cerebrovascular disease, arteriosclerosis, atherosclerosis, nonatheromatous arteriosclerosis, or vascular wall damage from cellular events leading toward immune mediated vascular damage, and inhibiting stroke or multiinfarct dementia.

15
20
25
30

The following examples are illustrative only and are not intended to be a limitation on the present invention.

EXAMPLES

5

Example 1 - General Procedures for Evaluation of Oxidative Impurities Present in CCI-779 Samples

The oxidative and hydrolytic impurities in a CCI-779 sample may be quantitated as a mixture of co-eluting materials over a specified range of retention times. The method used is a reverse phase gradient HPLC/UV procedure. The chromatographic conditions are outlined below.

10

Table 1 : Chromatographic Conditions (HPLV/UV) for Determination of Oxidative/Hydrolysis Degradation Products

Method Parameter	Requirement												
Column Specification:	Ultracarb™ ODS(30), 150 x 4.6 mm, 5 µm size particles												
Mobile Phase:	Mobile Phase A: 50% acetonitrile: 50% water Mobile Phase B: 80% acetonitrile: 20% water												
Gradient:	<table> <tr> <th>Time</th><th>% Mobile Phase B</th></tr> <tr> <td>0</td><td>0</td></tr> <tr> <td>40</td><td>100</td></tr> <tr> <td>60</td><td>100</td></tr> <tr> <td>60.1</td><td>0</td></tr> <tr> <td>75</td><td>0</td></tr> </table>	Time	% Mobile Phase B	0	0	40	100	60	100	60.1	0	75	0
Time	% Mobile Phase B												
0	0												
40	100												
60	100												
60.1	0												
75	0												
Column Temperature:	45°C												
Flow Rate:	1.5 mL/minute												
Detection:	UV, 225 nm												
Injection Volume:	50 µL												
Sample solvent	300 mL water: 700 mL acetonitrile: 0.5 mL acetic acid												
Sample Preparation:	2.0 mL of sample is transferred into a 25 mL volumetric flask and diluted to volume with sample solvent												
Calibration:	The amount of oxidation/hydrolysis degradation products is determined by comparing the appropriate peak areas in the sample preparation chromatograph to the total area of CCI-779 and its related compounds. The area between RRT ^a 0.13 to 0.81 is a measure of the												

	oxidation/hydrolysis degradation products. The reporting limit was 0.05%
	a. RRT= Relative retention time. Retention time of isomer B used as reference point
System Suitability Preparations	20 µg/mL of Temsirolimus reference standard in sample solvent. (solution A) 1 µg/mL of Temsirolimus reference standard in sample solvent. (solution B) 2 mg/mL of a Temsirolimus control batch in sample solvent.
<u>System Suitability</u>	
Sample solvent blank	No interfering peaks between 4-19 minutes
Signal to Noise (Solution B)	S/N >10
Retention time CCI-779 Isomer B (Solution A)	18 to 24 minutes
Theoretical Plates CCI-779 Isomer B (Solution A)	≥ 2000
Tailing Factor CCI-779 Isomer B (Solution A)	≤ 2.0
%CV (blank subtracted control samples)	≤ 10%
%RSD (6 standard injections):	≥ 100 ≤ 6.0%

Table 2

5

With reference to Table 1 above, the following table provides further details relating to the gradients used with mobile phase A (MP-A) and mobile phase B (MP-B). During "linear change", the gradient is changed at a uniform rate. During "isocratic hold", the solvent phase remains constant. During "step change", an

10 incremental change in the solvent is introduced.

Time (min)	% MP-A	% MP-B	Comments
0	100	0	Initial conditions
40	0	100	Linear change
60	0	100	Isocratic hold
60.1	100	0	Step change
75	100	0	Equilibrate for next injection

As an alternative to HPLC/UV, the CCI-779 in the sample was quantitated by analyzing the extent of one oxygen, two oxygen, 3 oxygen, one oxygen plus water, and water incorporation, based on the m/z of the addition product.

5

Table 3

Chromatographic Mass Spectrometric (HPLC/MS) Conditions for Determination of Oxidative/Hydrolysis Degradation Products

Method Parameter	Requirement																		
Column Specification	C18, 3 μm or 5 μm , 150 x 2.0 mm held at 45°C																		
Mobile Phase	<u>Mobile Phase A:</u> Mix 95 volumes of water with 5 volumes of acetonitrile and 0.1 volume of formic acid. <u>Mobile Phase B:</u> Mix 5 volumes of water with 95 volumes of acetonitrile and 0.1 volume of formic acid.																		
Gradient	<table><tr><td><u>Gradient:</u></td><td><u>Time(min)</u></td><td><u>%B</u></td></tr><tr><td></td><td>0</td><td>30</td></tr><tr><td></td><td>6</td><td>30</td></tr><tr><td></td><td>46</td><td>76</td></tr><tr><td></td><td>47</td><td>100</td></tr><tr><td></td><td>57</td><td>100</td></tr></table> Re-equilibrate at initial conditions for 13 minutes	<u>Gradient:</u>	<u>Time(min)</u>	<u>%B</u>		0	30		6	30		46	76		47	100		57	100
<u>Gradient:</u>	<u>Time(min)</u>	<u>%B</u>																	
	0	30																	
	6	30																	
	46	76																	
	47	100																	
	57	100																	
Flow Rate	0.2 mL/min																		
Detection	Mass spectrometric																		
Ionization Mode	Negative Electrospray																		
Single Ion Monitoring	m/z 1014.7, m/z 1044.7 , m/z 1046.71046, m/z 1060.7, m/z 1062.7 and m/z 1076.7																		
Sample Solvent	Acetonitrile (API) 30 :70 :0.5 (water: acetonitrile: acetic acid V:V:V) for Drug product																		
Standard	60 μg/mL, 50 μg/mL, 40 μg/mL, 30 μg/mL, 10 μg/mL, 4 μg/mL and 1 μg/mL of 1R,2R-Dihydroxyl Temsirolimus																		
Calibration	Calculate the quadratic regression line relating the peak areas (y) of the 1R,2R-Dihydroxyl Temsirolimus peak in the standard preparation chromatograms to the corresponding concentration value of the reference standard (x). Connect the low standard point to the origin.																		
Control Preparation	Temsirolimus preparation at a concentration of about 2 mg/ML																		

<u>System Suitability</u>	
Check Tune	Must pass for the range of m/z 800-1100
Retention time 1R, 2R, Dihydroxyl temsirolimus (30 µg/mL standard)	30 to 38 minutes
Theoretical Plates 1R, 2R, Dihydroxyl temsirolimus (30 µg/mL standard)	10,000
Tailing Factor 1R, 2R, Dihydroxyl temsirolimus (30 µg/mL standard)	≤2.0
Signal to noise (1 µg/mL standard)	≥10
R ² for the quadratic regression	≥0.990

Example 2 - Varying Levels of Impurities

In this example, three rapamycin compositions, each of which contains CCI-
 5 779 2.5%, d,l- α -tocopherol 0.075%, anhydrous citric acid 0.0025%, dehydrated
 alcohol 39.5%, and propylene glycol q.s., with varying levels of oxidative impurities,
 were monitored over a period of about 3 to 5 months to determine their stabilities at
 various temperatures and humidities. These batches contained about 0.5%, about 1%
 and about 2% of oxidative/hydrolysis impurities, respectively.

10 Aliquots of the formula were subdivided into glass vials, stoppered, sealed
 and stored at 5°C, 25°C/60% relative humidity (RH), or 40°C/75% RH. Samples
 were monitored for (i) appearance and description, (ii) moisture, (iii) strength, total
 related compounds (nonoxidative), (iv) oxidative/hydrolysis impurities, and (v) α -
 tocopherol content. The data illustrates that for samples initially containing about
 15 0.5% impurities, there was a slight increase in total (nonoxidative) degradation and
 oxidative/hydrolytic degradation after 1 month at 40°C. After 3 and 5 months at 25°C
 the formulation was stable, *i.e.*, the potency remained the same as did total
 degradation.

For the sample initially containing about 1% impurities, there was an increase
 20 in oxidative/hydrolytic degradation products to about 1.94% after three months. This
 trend continued out to 5 months at 25 °C, with increases in total nonoxidative and
 oxidative/hydrolytic degradation products to 1.65 and 2.3% respectively.

For the sample initially containing about 2% impurities, the total nonoxidative degradation and oxidative/hydrolytic degradation increased after 1 month 40 °C/75% RH to 8 and 4.3% respectively. After three months, both total nonoxidative degradation and oxidative/hydrolytic degradation products increased for the samples at 25°C/60%RH to 3.3 and 4.2% respectively. Figure 3 illustrates the effect of initial oxidative/hydrolysis degradant levels contributed by the input drug raw material on the stability of CCI-779 after 1 and 3 month storage.

In summary, it was found that higher initial concentrations of the oxidative/hydrolytic impurities adversely affected the stability of the CCI-779 samples. In fact, the greatest stability of samples containing CCI-779 occurred when the initial concentration of the oxidative impurities in the rapamycin composition was 0.5% or less. Therefore, reducing the initial oxidative impurities significantly enhances the shelf-life of the formulated CCI-779 product.

Example 3 - Varying α -Tocopherol Concentration

To further investigate the effect of the oxidative impurities in rapamycin compositions, studies were conducted by varying concentrations of α -tocopherol in the rapamycin compositions.

Samples of CCI-779 containing 0.2%, 0.5% and 1% d,l- α -tocopherol (Eisai) were placed in 2 mL flint glass vials and stoppered with 13 mm West Teflon Faced 4432/50 stoppers. The effect of increased α -tocopherol concentrations on the rapamycin compositions was monitored over 1 month at 40°C. Samples were stored upright at about 5°C or about 40°C.

After 1 month at 40°C, the data illustrate that in all of the samples, the α -tocopherol concentration dropped significantly. However, for the samples containing 0.2% and 0.5% of α -tocopherol, the concentration of the oxidative impurities remained essentially unchanged, *i.e.*, the concentration of oxidative impurities did not increase. However, there was an overall loss of potency of the samples due to the formation of other degradation products.

For samples containing 1% α -tocopherol, the presence of the oxidative impurities drastically increased to 8.42%.

In summary, increasing the concentration of α -tocopherol in the samples to 0.2 and 0.5% slowed the growth of oxidative impurities, but did not inhibit degradation of the CCI-779 via other mechanisms when the oxidative impurity levels were 3% or more. As indicated in the previous example, inhibition of the non-oxidative
5 impurities was controlled by limiting the initial amount of oxidative/hydrolysis impurities in the drug substance.

All documents listed in this specification are incorporated herein by reference. While the invention has been described with reference to particular embodiments, it
10 will be appreciated that modifications can be made without departing from the spirit of the invention. Such modifications are intended to fall within the scope of the appended claims.

WHAT IS CLAIMED IS:

1. A method of preparing a rapamycin composition having increased potency, said method comprising the steps of:
selecting a rapamycin compound having less than 1.5% oxidative and hydrolytic rapamycin impurities; and
formulating the selected rapamycin with an antioxidant and optional excipients.
2. The method according to claim 1, wherein the selecting step comprises screening the rapamycin in a high performance liquid chromatography assay.
3. The method according to claim 1 or 2, wherein the antioxidant is selected from the group consisting of a tocopherol, vitamin C, 2,6-di-tert-butyl-4-methylphenol, and mixtures thereof.
4. The method according to claim 3, wherein the antioxidant is α -tocopherol.
5. The method according to any one of claims 1 to 4, wherein the selected rapamycin has less than 0.5% oxidative impurities.
6. The method according to any one of claims 1 to 5, wherein the selected rapamycin is formulated for parenteral delivery.
7. The method according to any one of claims 1 to 6, wherein the selected rapamycin is formulated as a liquid concentrate.
8. The method according to claim 7, wherein the selected rapamycin is formulated with d,l- α -tocopherol, anhydrous citric acid, dehydrated alcohol, and propylene glycol.

9. The method according to any one of claims 1 to 5, wherein the selected rapamycin is formulated for oral delivery.

10. A method of preparing a rapamycin composition having increased potency, said method comprising the steps of:

selecting a rapamycin compound having less than 1.5% oxidative and hydrolytic rapamycin impurities;


formulating the selected rapamycin with at least two antioxidants and optional excipients.

11. The method according to claim 10, wherein at least one of the antioxidants is vitamin C or 2,6-di-tert-butyl-4-methylphenol.

12. The method according to claim 10, wherein said at least two antioxidants are vitamin C and 2,6-di-tert-butyl-4-methylphenol.

13. The method according to any one of claims 1 to 12, wherein said rapamycin is selected from the group consisting of rapamycin and CCI-779.

FIG. 1







FIG. 2

FIG. 3



INTERNATIONAL SEARCH REPORT

International application No

PCT/US2006/048329

A. CLASSIFICATION OF SUBJECT MATTER

INV. A61K31/05 A61K31/355 A61K31/375 A61K31/436

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, BIOSIS, EMBASE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	AU 759 219 B2 (NOVARTIS AG [CH]) 10 April 2003 (2003-04-10) claims 1-4 examples 1,2 -----	1-13
X	US 2003/191148 A1 (NAVARRO FRANCOIS [FR] ET AL) 9 October 2003 (2003-10-09) paragraphs [0005], [0006], [0012], [0018], [0019], [0128] claims 1-8 -----	1-13
X	WO 2005/010010 A (WYETH CORP [US]; ZHU TIANMIN [US]; MATTES JAMES F [US]) 3 February 2005 (2005-02-03) page 10, line 22 - line 31 ----- -/-	1-13

☒ Further documents are listed in the continuation of Box C.☒ See patent family annex.

* Special categories of cited documents :

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

29 May 2007

Date of mailing of the international search report

06/06/2007

Name and mailing address of the ISA/

European Patent Office, P.B. 5816 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Albayrak, Timur

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2006/048329

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 329 460 A2 (AMERICAN CYANAMID CO [US]) 23 August 1989 (1989-08-23) page 3, line 32 - line 41	1-13
X	WO 2004/011000 A (WYETH CORP [US]; RUBINO JOSEPH T [US]; SISKAVICH VICTORIA [US]; HARRIS) 5 February 2004 (2004-02-05) page 3 claims 1-30	1-13

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2006/048329

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
AU 759219	B2	10-04-2003	
		AR 048073 A2	29-03-2006
		AR 026102 A1	29-01-2003
		AU 1657300 A	26-06-2000
		BE 1012869 A3	03-04-2001
		BR 9915986 A	04-09-2001
		CA 2351580 A1	15-06-2000
		CN 1374872 A	16-10-2002
		CZ 20012001 A3	12-09-2001
		WO 0033878 A2	15-06-2000
		EP 1137439 A2	04-10-2001
		FR 2786771 A1	09-06-2000
		HU 0104489 A2	28-03-2002
		ID 29250 A	16-08-2001
		IT MI992520 A1	04-06-2001
		JP 3805625 B2	02-08-2006
		JP 2002531527 T	24-09-2002
		JP 2005200429 A	28-07-2005
		JP 2006111637 A	27-04-2006
		KR 20060096477 A	11-09-2006
		NO 20012424 A	16-05-2001
		NZ 511936 A	31-10-2003
		NZ 527781 A	26-11-2004
		PL 348333 A1	20-05-2002
		SK 7662001 A3	03-12-2001
		TR 200101416 T2	21-01-2002
		TR 200201428 T2	23-12-2002
		TW 248938 B	11-02-2006
		US 2002032213 A1	14-03-2002
US 2003191148	A1	09-10-2003	NONE
WO 2005010010	A	03-02-2005	
		AU 2004259669 A1	03-02-2005
		BR PI0412659 A	26-09-2006
		CA 2528173 A1	03-02-2005
		CN 1823073 A	23-08-2006
		EP 1646635 A1	19-04-2006
		MX PA06000407 A	17-03-2006
EP 0329460	A2	23-08-1989	
		AU 3006589 A	24-08-1989
		BR 8900688 A	10-10-1989
		CA 1323835 C	02-11-1993
		DD 297821 A5	23-01-1992
		DE 68901543 D1	25-06-1992
		DK 75689 A	19-08-1989
		ES 2031351 T3	01-12-1992
		GR 3004648 T3	28-04-1993
		HU 50629 A2	28-03-1990
		IE 61053 B1	21-09-1994
		JP 2003690 A	09-01-1990
		JP 2843591 B2	06-01-1999
		NZ 228038 A	26-07-1991
		PT 89760 A	04-10-1989
		ZA 8901248 A	25-10-1989
WO 2004011000	A	05-02-2004	
		AU 2003254168 A1	16-02-2004
		BR 0313024 A	12-07-2005
		CA 2493878 A1	05-02-2004
		CN 1671385 A	21-09-2005

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2006/048329

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
---	---------------------	----------------------------	---------------------

WO 2004011000 A

EP	1553940 A1	20-07-2005
JP	2005537285 T	08-12-2005
KR	20050026025 A	14-03-2005
MX	PA05001013 A	16-05-2005
NZ	537829 A	29-09-2006